

Total No. of Printed Pages—7

**5 SEM TDC MTMH (CBCS) C 11**

**2024**

( November )

**MATHEMATICS**

( Core )

Paper : C-11

( **Multivariate Calculus** )

*Full Marks : 80*

*Pass Marks : 32*

*Time : 3 hours*

*The figures in the margin indicate full marks  
for the questions*

1. (a) State the range of

$$f(x, y) = \frac{1}{xy}$$

1

- (b) Fill in the blank :

1

The definition of limit of  $f(x, y)$  applies to the boundary points and \_\_\_\_\_ of the domain of  $f$ .

( 2 )

(c) State when

$$\lim_{(x, y) \rightarrow (x_0, y_0)} f(x, y)$$

does not exist.

1

(d) Investigate the existence of the limit

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{x^2 + y}{y}$$

2

(e) Find  $f_x$  and  $f_y$ , given  $f(x, y) = \log_y x$ .

2

(f) Use chain rule to find  $\frac{dw}{dt}$  where  $w = x^2 + y^2$  along the path  $x = \cos t$  and  $y = \sin t$  at  $t = \pi$ .

2

(g) If  $w = f(x, y)$  is differentiable and both  $x$  and  $y$  are differentiable of  $t$ , then show that  $w$  is a differentiable function of  $t$  and

$$\frac{dw}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

3

Or

If  $w = yz + zx + xy$  and  $x = r + s$ ,  
 $y = r - s$ ,  $z = rs$ , then find  $\frac{\partial w}{\partial r}$  and  $\frac{\partial w}{\partial s}$   
at  $(r, s) = (2, 1)$ .

( 3 )

(h) Determine  $\nabla f$  at  $(1, 2, -2)$  where

$$f(x, y, z) = (x^2 + y^2 + z^2)^{-\frac{1}{2}} + \log xyz$$

4

(i) Find the directional derivative of

$$f(x, y, z) = 3e^x \cos yz$$

at the origin in the direction of

$$2\hat{i} + \hat{j} - 2\hat{k}$$

4

Or

Find the tangent plane and normal to the surface  $x + y + z = 1$  at  $(0, 1, 0)$ .

(j) Find the local extrema or saddle point as applicable of the function

$$f(x, y) = 6x^2 - 2x^3 + 3y^2 + 6xy$$

5

(k) Use the method of Lagrange's multiplier to maximise  $f(x, y) = xy$  subject to the constraint  $x + y = 16$ .

5

Or

Find the points on the surface  $z^2 = xy + 4$  closest to the origin.

( 4 )

2. (a) Find the  $y$ -limits of integration for the integral

$$\iint_R f(x, y) dA$$

where  $R$  is the region bounded by the line  $x+y=1$  and a circle of radius 1 with its centre at the origin.

1

- (b) Sketch the region of integration on the plain paper for the integral

$$\int_0^3 \int_0^2 f(x, y) dx dy$$

1

- (c) Write any one iterated integral for the double integral

$$\iint_R f(x, y) dA$$

where  $R$  is the triangular region with the vertices  $(0, 0)$ ,  $(1, 0)$  and  $(0, 1)$ .

2

- (d) Reverse the order of integration

$$\int_0^1 \int_2^{4-2x} dy dx$$

2

- (e) Find the volume of the region between the cylinder  $z=y^2$  and  $xy$ -plane bounded by the planes  $x=0$ ,  $x=1$ ,  $y=-1$ ,  $y=1$ .

4

( 5 )

- (f) Set up the iterated integral for evaluating the integral

$$\iiint_D f(r, \theta, z) dz r dr d\theta$$

over the region  $D$  which is a prism with its base on the  $xy$ -plane bounded by the  $x$ -axis and the lines  $y=x$  and  $x=1$  and whose top lies in the plane  $z=2-y$ .

5

- (g) Write any two different iterated triple integrals for determining the volume of the tetrahedron bounded by the coordinate planes and the plane

$$x + \frac{y}{2} + \frac{z}{3} = 1 \qquad 2\frac{1}{2} + 2\frac{1}{2} = 5$$

3. (a) Define Jacobian of the transformation

$$x = f(u, v); \quad y = g(u, v)$$

1

- (b) If

$$u = \frac{1}{2}(x+1), \quad v = \frac{1}{3}(y+4); \quad w = 2z+4$$

find  $\frac{\partial(x, y, z)}{\partial(u, v, w)}$ .

3

( 6 )

2.

- (c) Evaluate  $\int_C (xy + y + z) ds$  along the curve

$$\vec{r}(t) = 2t\hat{i} + t\hat{j} + (2-2t)\hat{k}; 0 \leq t \leq 1 \quad 3$$

- (d) Show that the work done  $\int \vec{F} \cdot d\vec{r}$  around every closed loop in an open region  $D$  is zero if and only if  $\vec{F}$  is conservative in  $D$ . 4

- (e) State the fundamental theorem on line integrals. If  $\vec{F}$  is a vector field whose components are continuous throughout an open connected region  $D$  in space, then there exists a differentiable function  $f(x, y, z)$  such that  $\vec{F} = \nabla f$ , show that

$$\int_A^B \vec{F} \cdot d\vec{r}$$

is path independent where

$$\vec{r} = g(t)\hat{i} + h(t)\hat{j} + k(t)\hat{k}; a \leq t \leq b$$

is a smooth curve joining  $A$  and  $B$  in  $D$ .

$$1+3=4$$

4. (a) Find the curl of the function

$$\vec{F}(x, y) = (x^2 - y)\hat{i} + (xy - y^2)\hat{j} \quad 1$$

- (b) State Green's theorem in circulation-curl or tangential form. 2

P25

P25/171

( Continued )

( 7 )

- (c) Integrate  $f(x, y, z) = x + y + z$  over the surface of the cube cut from the first octant by the planes  $x = a; y = a; z = a$ . 3

- (d) Evaluate  $\iint_S f d\sigma$  where  $S$  is the surface area of the cone  $z = \sqrt{x^2 + y^2}; 0 \leq z \leq 1$  and  $f(x, y, z) = x^2$ . 4

- (e) State and prove Stokes' theorem. 5

Or

State and prove divergence theorem.

\*\*\*

P25—2000/171

5 SEM TDC MTMH (CBCS) C 11

