

Total No. of Printed Pages—7

5 SEM TDC PHYH (CBCS) C 11

2025

(Nov/Dec)

PHYSICS

(Core)

Paper : C-11

(Quantum Mechanics and Applications)

Full Marks : 53

Pass Marks : 21

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

1. Choose the correct answer :

1×5=5

(a) Quantum mechanical operator of total energy is

(i) $\frac{\hbar}{i} \nabla$

(ii) $i\hbar \frac{\partial}{\partial t}$

(iii) $\frac{\hbar}{i} (r \times \nabla)$

(iv) $-\frac{\hbar^2}{2m} \nabla^2$

(2)

(b) The energy of a linear harmonic oscillator is

- (i) discrete and equispaced
- (ii) continuous
- (iii) partly discrete and partly continuous
- (iv) discrete but not equispaced

(c) If A is the operator, operating on a state function ψ , then the possible values of any physical quantity of a system in the operator equation $A\psi = a\psi$ are given by

- (i) eigenfunction ψ
- (ii) eigenvalue ψ
- (iii) eigenvalue a as well as eigenfunction ψ
- (iv) None of the above

(d) In Schrödinger picture, time evolution operator is

- (i) time dependent
- (ii) time independent
- (iii) constant
- (iv) None of the above

26P/478

(Continued)

(3)

(e) The selection rule for radiative transition of an electron in hydrogen atom is

- (i) $\Delta l = 0, \Delta m_l = \pm 1$
- (ii) $\Delta l = \pm 1, \Delta m_l = \pm 1$ or 0
- (iii) $\Delta l = \pm 1, \Delta m_l = 0$
- (iv) $\Delta l = \pm 1, \Delta m_l = \pm 1$

2. (a) What do you understand by quantum mechanical operators? 2

(b) Normalise $\psi(x) = Ae^{-\alpha x^2}$ over the interval $-\infty \leq x \leq +\infty$. 2

(c) Define Larmor frequency. 2

Or

What are symmetric and anti-symmetric wave functions?

(d) Show that the expectation value of Hamiltonian operator (H) is the total energy (E) of the system. 2

3. (a) What is Stark effect? How many components are observed in Stark effect? 1+2=3

26P/478

(Turn Over)

(4)

Or

Find the magnetic moment, in Bohr magneton, of an atom in the 3P_2 state. 3

- (b) The ground state of a harmonic oscillator is given by $\psi(x) = Ae^{-\alpha^2 x^2}$ where $\alpha^2 = \frac{k}{2\hbar\omega}$, k is the quasistatic force constant ($U = \frac{1}{2}kx^2$). Calculate the mean values of the kinetic and potential energies of the oscillator. 3

Or

Show that for large quantum numbers, the classical probability and quantum mechanical probability of an linear harmonic oscillator become identical. 3

- (c) Derive one-dimensional Schrödinger equation from a wave packet and hence show that the energy operator and momentum operator can be represented as follows : 3

$$E \rightarrow i\hbar \frac{\partial}{\partial t} \text{ and } P \rightarrow i\hbar \frac{\partial}{\partial x}$$

(5)

- (d) Prove that the relation

$$\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot \vec{J} = 0$$

where $\vec{J}$ is the probability current density and ρ is the probability density. 3

Or

State and prove Heisenberg's uncertainty principle for wave packets. 3

4. (a) How do you represent the states of an atom in spectral notation? 4

Or

Show that if $\psi_1(\vec{r})$ and $\psi_2(\vec{r})$ are the independent solutions of the Schrödinger equation, then $\psi(\vec{r}) = \alpha_1\psi_1(\vec{r}) + \alpha_2\psi_2(\vec{r})$ is also a solution. What does it imply? 3+1=4

- (b) Explain L-S coupling and draw the vector diagram of LS coupling. Under what condition LS coupling breaks down? 4

(6)

5. (a) Describe Stern-Gerlach experiment. Discuss how it explained space quantization and electron spin. 5

- (b) Write down the Schrödinger wave equation for the hydrogen atom. Separate the Schrödinger equation for the motion of an electron in hydrogen atom into one radial and two angular parts. 5

Or

Show that the orbital angular momentum of the electron in hydrogen atom is $L = \hbar\sqrt{l(l+1)}$, where l is the orbital quantum number. 5

- (c) Show that the energy eigenvalues of a particle in a one-dimensional box of infinite depth are discrete and proportional to n^2 . Calculate the value of lowest energy of an electron in a one-dimensional force-free region of length 4 Å. 4+1=5

- (d) Find the values of $\langle x \rangle$, $\langle x^2 \rangle$, $\langle p_x \rangle$ and $\langle p_x^2 \rangle$ for the Gaussian wave packet

$$\psi(x) = \left(\frac{1}{\sigma\sqrt{\pi}} \right)^{\frac{1}{2}} e^{-\frac{x^2}{2\sigma^2}} e^{ik_0x}$$

5

(7)

Or

The wave function of a particle moving in one-dimension is given by

$$\psi(x) = \begin{cases} \sqrt{\frac{15}{\alpha}} A(\alpha^2 - x^2), & \text{for } -\alpha \leq x \leq \alpha \\ 0, & \text{for } |x| > \alpha \end{cases}$$

- (i) Find the value of A that will normalise $\psi(x)$.
 (ii) Calculate the expectation value of x and p . 3+2=5
